

The Role of Host Genetic Variability in the Development and Establishment of Human Gut Microbiome Diversity

Larissa Matos

Genome USP, Brazil

Abstract

Background: Studies have shown that human microbiome plays important roles in physiology, from food digestion to mental diseases. Since the gut microbiome composes the highest number of microbial cells outnumbering even our own cell counts, it is expected that the gut microbiome would affect a great deal of human biological functions. This makes the gut microbiome key to maintain homeostasis in the various biological levels where there are constant and active interactions between microbes, tissue, cells and molecules. The structure of the gut microbiota is shaped by many factors, including host genetics. Understanding how these factors determine the microbiome during the development and establishment of the gut microbiota at early stages of human life is crucial to infer biological and pathological microbiome composition. The purpose of this study is to apply tools of bioinformatics to process and analyse data samples of feces of triplet babies in order to verify host genetic associations with gut microbiome during the first 3 years of the babies lives.

Methods: Infant feces are collected from the first week of life until 3 years of age. DNA extraction from the samples is performed followed by PCR (Polymerase Chain Reaction) targeting the specific bacterial 16S ribosomal RNA gene, region V3 - V4. Next-Generation sequencing is applied. Amplicons are then computationally processed by efficient algorithms to yield high quality reads. Later, these reads are assigned to groups of taxa based on differences on single nucleotide. Species specific classifiers and database are used for taxonomic assignment. Alpha and beta diversity are analysed with both qiime2 plugins and R packages.

Results: Most of our samples presented enough reads to identify all taxa. Phylogenetic diversity increased in samples on later time points presenting time as the dominant factor to determine alpha and beta diversity. Actinobacteria, Verrucomicrobia, Proteobacteria, Firmicutes and Bacteroidetes were the dominant phyla in all samples. At the species level, monozygotic twins presented more similar microbiome between them than between their dizygotic sibling.

Conclusions: Our results showed that genetic factors could be detected at the species level and that time is crucial for diversity in the first months of life. The identification of the gut microbiome structure in both monozygotic and dizygotic twins sheds some light on how the development and establishment of microbiota take place on the human gut. For our next analysis we will apply appropriate statistics for twin studies and check specific species and its association with host genetics.

Biography

Larissa Matos graduated in biomedicine (2010) and master in biology applied to health (2013) from the Federal University of Pernambuco - UFPE. She is PhD in science (2017) from the University of São Paulo - USP (Program of Experimental Epidemiology Applied to Zoonoses). She has experience in the areas of microbiology, molecular biology, carcinogenesis and

molecular probes, with a focus on microbiome, bacterial pathogenesis (leptospirosis), NGS sequencing, bioinformatics, cloning, expression and purification of recombinant proteins, electrophoresis, chromatography, ELISA, western blotting, chemiluminescence and diagnosis. She is currently a postdoc researcher at the Center for Research on the Human Genome and Stem Cells (CEGH-IB/USP), working mainly with the microbiome-host interaction.